




2023 edition

## Immunization Schedule

Name of Vaccine	Administered	0year	1month	2months	3months	4months	5months	6months	7months	8months	9-11months	12-15months	16-17months	18-23months	2Age	3Age	4Age	5Age	6Age	7Age	8Age	9Age	10Age
Haemophilus influenzae type b (Hib)	Inactivated vaccine			①	②	③	If the child did not receive their first vaccination before 7 months after birth, then the number of vaccines that should be given is different. It is important to pay attention to the age the first vaccine was given.					④	The 4th vaccination should be given over 1 year old.										
Pneumococcal conjugate	Inactivated vaccine			①	②	③	every 4 weeks					④											
Hepatitis B (excluding prevention of perinatal infection)	Universal	Inactivated vaccine		①	②	20 weeks (about 5 months)				③	The 3rd vaccination should be given at an interval of about 5 months after the 1st vaccination.												
	Maternal and child infection prevention	Inactivated vaccine	①	②	The 1st vaccination should be given by 6 days and 14 weeks after birth				③														
Rotarix	Monovalent	Live vaccine		①	②	every 4 weeks					You should pick just one Rotavirus vaccine. You should complete the same vaccine you picked.												
RptaTeg	5-valent			①	②	③	If the vaccine is given in intervals of 4 weeks between vaccinations, then these vaccines can be administered at the same time: Rota, HepB, Hib, Pneumococcal conjugate and diphtheria, tetanus, pertussis & inactivated polio					④											
Diphtheria, tetanus, pertussis, & inactivated polio DPT/IPV	Inactivated vaccine	At least 4 times vaccinations should be given.		①	②	③	every over 3 weeks					④	The 4th vaccination should be given at an interval of at least 6 months after the 3rd vaccination. The best interval is after 1 year after 3rd vaccination.										
DPT+IPV	Inactivated vaccine																						
Diphtheria, tetanus DT	Inactivated vaccine																						
BCG	Live vaccine	Don't be given at under 3 months after birth.		①	The best period for BCG is when your child is over 5 months after birth.																		
MR (measles and rubella combined)	Live vaccine											①											
Varicella (chicken pox)	Live vaccine											①	②	The 2nd vaccination should be given at an interval of over 3 months after the 1st vaccination.									
Mumps	Live vaccine											①	②	The 2nd vaccination is recommended to be administered simultaneously with MR.									
Japanese encephalitis	Inactivated vaccine															①	③	from 3 years old.			④	9~12age	
Influenza	Inactivated vaccine																						
HPV (Human papillomavirus)	Inactivated vaccine																						

Recommended time period for routine vaccination

Target age for routine vaccination

Recommended time period for voluntary vaccination

Age when voluntary vaccines may be administered

Time period for vaccination by health insurance



Uttaro Genki note

Let's save your family's immunization records and share them with doctors. You can reserve vaccinations online. <https://e-chusya.com/>